

Supplemental Table 2: Baseline and Hospitalization Characteristics of Children with GSW by Initial Disposition

Characteristic	Admitted (n = 200)	Not Admitted*		p-value
		Died** (n=14)	Discharged (n =79)	
Age _{continuous} , mean (sd)	13.10 (3.9)	13.05 (3.51)	14.23 (2.26)	0.055
Age _{cat-SIPA} , n (%)				0.051
0-6	24 (12.0)	1 (7.1)	2 (2.5)	
7-12	23 (11.5)	4 (28.6)	10 (12.7)	
13-18***	153 (76.5)	9 (64.3)	67 (84.8)	
Female, n (%)	38 (19.0)	3 (21.4)	17 (21.5)	0.882
Race, n (%)				0.918
White	3 (1.5)	0 (0.0)	1 (1.3)	
Black or African American	185 (92.5)	12 (85.7)	72 (91.1)	
More than one race	5 (2.5)	1 (7.1)	2 (2.5)	
Unknown / not reported	7 (3.5)	1 (7.1)	4 (5.1)	
Ethnicity, n (%)				0.971
Hispanic or Latino	12 (6.0)	1 (7.1)	4 (5.1)	
NOT Hispanic or Latino	178 (89.0)	12 (85.7)	72 (91.1)	
Unknown / not reported	10 (5.0)	1 (7.1)	3 (3.8)	
Insurance Status, n (%)				0.117
Public	126 (63.0)	6 (42.9)	43 (54.4)	
Private	18 (9.0)	0 (0.0)	6 (7.6)	
Uninsured	56 (28.0)	8 (57.1)	30 (38.0)	
SIPA ^a , n (%)	50 (25.0)	3 (21.4)	14 (17.7)	0.424
ISS, mean (sd)	11.90 (13.8)	44.43 (32.20)	1.62 (2.68)	<0.001
GCS _{cat} : < /=8, n (%)	23 (11.5)	10 (71.4)	0 (0.0)	<0.001
Surgery, n (%)	114 (57.0)	6 (42.9)	-	-
Ventilation, n (%)	50 (25.0)	-	-	-
Mortality, n (%)	10 (5.0)	14 (100.0)	0 (0.0)	<0.001
PICU Admission, n (%)	120 (60.0)	-	-	-
Hospital LOS in days, median (IQR)	3.00 (2.0-8.0)	-	-	-

* One patient was transferred from the pediatric to the adult hospital and not included among discharged patients.

**Deaths occurred in the Emergency Department or the operating room.

***One patient in the non-admitted group was 18 years old. All other patients were < 17 years old.

^aCut-off Shock Index Values for significant injury by age: >1.22 (age 0–6), >1.0 (7–12), >0.9 (13–16)

SIPA = Shock Index Pediatric Adjusted, ISS = Injury Severity Score, GCS = Glasgow Coma Scale, LOS= length of stay